

# Avocado and Cardiovascular Health

Camila Weschenfelder<sup>1</sup>, Júlia Lorenzon dos Santos<sup>1</sup>, Priscilla Azambuja Lopes de Souza<sup>1</sup>, Viviane Paiva de Campos<sup>1</sup>, Aline Marcadenti<sup>1,2\*</sup>

<sup>1</sup>Postgraduate Studies Program in Cardiology, Instituto de Cardiologia/Fundação Universitária de Cardiologia do Rio Grande do Sul (IC/FUC), Porto Alegre, Brazil

<sup>2</sup>Department of Nutrition, Federal University of Health Sciences of Porto Alegre (UFCSPA), Porto Alegre, Brazil  
Email: \*[marcadenti@yahoo.com.br](mailto:marcadenti@yahoo.com.br)

Received 30 June 2015; accepted 25 July 2015; published 29 July 2015

Copyright © 2015 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

---

## Abstract

Avocado is a fruit which had a caloric density of 1.7 kcal per gram and a half unit (~70 g) is composed by 114 kcal, 4.6 g of fibers, 345 mg of potassium, 19.5 mg of magnesium, 1.3 mg of vitamin E and 57 mg of phytosterols. Approximately 75% of fiber's avocado contents are considered insoluble and 25% are soluble. The avocado contains lipids that consist of 71% from monounsaturated fatty acids (MUFA), 13% from polyunsaturated (PUFA) and 16% from saturated fatty acids (SFA). Recent researches have shown that avocado may improve hypercholesterolemia and may be useful in the treatment of hypertension and type 2 diabetes mellitus (T2DM). This way, avocado plays an important role in the cardiovascular health. This review summarizes the potential benefits of avocado consumption in the prevention of cardiovascular risk factors and metabolic diseases.

## Keywords

Persea, Diabetes Mellitus, Type 2, Dyslipidemias, Nutritional Status, Blood Pressure

---

## 1. Introduction

The tree of the avocado is originally from Mexico and Central America, and belongs to the *Lauraceae* family, genus *Persea* [1]-[3]. This genus is divided into two subgenera: *Persea* and *Eriodaphne* [3]. However, there are two important species in horticulture: *Persea americana* Mill and *Persea drymifolia* Cham.; this last comprising Mexican race avocados, which currently is considered as the botanical variety of *Persea americana* [3] [4].

The avocado is a fruit which had a caloric density of 1.7 kcal per gram and a half unit (~70g) is composed by 114 kcal, 4.6 g of fibers, 345 mg of potassium, 19.5 mg of magnesium, 1.3 mg of vitamin E and 57 mg of phy-

---

\*Corresponding author.

tosterols (**Table 1**) [5]. Approximately 75% of fiber's avocado contents are considered insoluble and 25% are soluble [6]. The avocado contains lipids that consist of 71% from monounsaturated fatty acids (MUFA), 13% from polyunsaturated (PUFA) and 16% from saturated fatty acids (SFA) [7]. Its high MUFA content in a water-based matrix appears to enhance the bioavailability of nutrients and phytochemical compounds of the avocado [8].

The avocado consumption has been related to benefits on some cardiovascular risk factors [5] [9], as well as to the improvement of the dietary quality and nutrient intake by increasing the consumption of MUFA, dietary fiber, magnesium, potassium, vitamins E and K [10]. Recent researches have shown that avocado can improve hypercholesterolemia and be useful in the treatment of hypertension (HTN) and type 2 diabetes mellitus (T2DM). This way, avocado can play an important role in the cardiovascular health.

This review aims to summarize the potential benefits of avocado consumption in the prevention of cardiovascular risk factors and metabolic diseases.

### 1.1. Avocado and Type 2 Diabetes Mellitus

The World Health Organization (WHO) have update the prevalence of T2DM in the United States (estimated about 30.3 million people by 2025) and worldwide, which is estimated around 380 million people diagnosed until the same year [11]. T2DM is associated with obesity, unhealthy diet, sedentary lifestyle and aging population

**Table 1.** Nutritional composition of the Hass avocado (*Persea americana*)<sup>5</sup>.

Nutrient/Phytochemical	Value per 100 g
<b>Proximates</b>	
Water (g)	72.3
Energy (Kcal)	167
Protein (g)	1.96
Fat (g)	15.4
Carbohydrate (g)	8.64
Fiber, total dietary (g)	6.8
Sugars, total (g)	0.3
<b>Minerals</b>	
Magnesium (mg)	29
Phosphorus (mg)	54
Potassium (mg)	507
Sodium (mg)	8
Zinc (mg)	0.68
Selenium (ug)	0.4
<b>Vitamins/Phytochemicals</b>	
Vitamin C (mg)	8.8
Folate, food (μg)	89
Lutein + zeaxanthin (μg)	271
Vitamin E (alpha-tocopherol) (mg)	1.97
<b>Lipids</b>	
Fatty acids, total saturated (g)	2.13
Fatty acids, total monounsaturated (g)	9.8
Fatty acids, total polyunsaturated (g)	1.82
Cholesterol (mg)	0

Adapted from Dreher ML *et al.*<sup>5</sup>.

[12]. Patients with fasting plasma glucose  $\geq 126$  mg/dL, oral glucose tolerance test (75 g glucose load)  $\geq 200$  mg/dL or glycated hemoglobin  $\geq 6.5\%$  [13] are diagnosed with T2DM. Patients with diabetes are at high risk for microvascular (*e.g.*, nephropathy, retinopathy and neuropathy) and macrovascular complications (*e.g.*, peripheral vascular disease, stroke and cardiovascular disease) [12].

Avocado has low sugar content (0.2 g in a half unit). D-mannoheptulose is the main kind of sugar found in the fruit but it seems to not have nutritional properties, appearing to be one more phytochemical component of the avocado [9]. The aqueous extract from the avocado's seeds has hypoglycemic agents, which act protecting against toxicity and oxidative stress [14]-[16]. In rats, phenolic extracts of avocado (from leaves and fruits) inhibited the activity of enzymes related to the development of T2DM ( $\alpha$ -amylase and  $\alpha$ -glucosidase), as well as the malondialdehyde production (MDA), a marker of oxidative stress and responsible for increasing the lipid peroxidation [17]. The hypoglycemic effect of the avocado was also related to its ability to stimulate the remaining pancreatic  $\beta$ -cells in animal models, making them able to secrete more insulin [18].

Diets rich in MUFA are considered alternatives for the dietary treatment of T2DM [19] and since avocado has a substantial amount of MUFA it could be used as an option for glycemic control in diabetic patients. However, few studies have evaluated the use of avocado in individuals with T2DM. Among overweight and moderately obese individuals, adding half avocado (70 g) in the lunch increased the satiety in a period of 3 to 5 subsequent hours, followed by a reduction of the insulin secretion in a 3-hour postprandial period [20]. Patients with hypercholesterolemia and T2DM supplemented with 300g/day of avocado for 7 days had their total cholesterol (TC) and LDL-cholesterol decreased by 17% and 22% respectively, and their triglycerides (TG) levels reduced by 22%; there was also a slightly increase in HDL-cholesterol when compared to the control group (isocaloric diet, 50% of total calories from fats and without avocado) [21].

The avocado paste can be obtained by the fruit oil and its effects were evaluated in rats, who consumed a hypercholesterolemic diet added of glucose solution and also the paste of avocado. Authors concluded that the animals had lower levels of blood sugar, lower values of the Homeostasis Model Assessment-Insulin Resistance Index (HOMA-IR Index) and less accumulation of fat in their liver. In this study, the improvement of the HOMA-IR Index and of the hepatic steatosis was attributed to the phytochemicals components and dietary fibers of the avocado [22].

## 1.2. Avocado and Dyslipidemia

Dyslipidemia is defined as lipid metabolic changes resulting from disturbances in any phase of the lipid metabolism, which impact on serum lipoproteins levels. It is an important cardiovascular risk factor: about a third of ischemic heart diseases are attributable to increased levels of TC. Globally, higher levels of cholesterol are responsible for 2.6 million deaths annually, and the treatment of dyslipidemia may reduce the cardiovascular risk by 30% over a period of 5 years [23] [24].

Nutritional therapy and changes in lifestyle are part of the non-pharmacological treatment for dyslipidemia. The American Heart Association/American College of Cardiology (AHA/ACC) recommends a healthy eating pattern, with 5% to 6% of total daily calories from SFA to reduce the levels of LDL-cholesterol [25]; SFA should be replaced by MUFA and PUFA intake [26] [27]. MUFA have a beneficial effect on hypercholesterolemia without increasing the lipid oxidation, in contrast with excessive intake of PUFA; MUFA ingestion also does not reduce serum HDL-cholesterol levels [28]. Recently, the PREDIMED study (*Prevención con Dieta Mediterránea*) showed that the Mediterranean diet supplemented with foods rich in MUFA (olive oil and nuts) reduced the incidence of major cardiovascular events by 30% after a follow-up of 4.8 years, in subjects at high risk for cardiovascular disease [29].

The lipid-lowering effect of avocado (also rich in MUFA) occurs mainly due to its phytosterol  $\beta$ -sitosterol [30]. Among 17,567 participants of the *National Health and Nutrition Examination Survey* (NHANES: 2001-2008) who had their avocado intake evaluated, the average daily intake was about a half of unit ( $70.1 \pm 5.4$ g/day), and the avocado consumers had higher levels of HDL-cholesterol when compared to those who did not consume [10].

The first clinical trial evaluating the influence of the avocado on serum TC was carried out by C. Grant Wilson in 1960 [31]. At that time, 16 men aged between 27 and 71 years (with and without hypercholesterolemia) were advised to consume 0.5 to 1.5 units of avocado a day. After 4 weeks, 8 participants had their TC reduced by 8.7% to 42.8%, without changing other lipid parameters. Subsequently, Carranza *et al.* [32] evaluated the effect of two diets in 16 individuals with dyslipidemia: 1) a diet rich in avocado (75% of total fat); or 2) a low sa-

turated fat/low dietary cholesterol diet. After 4 weeks, results showed that individuals allocated to the avocado diet had lower levels of TC and LDL-cholesterol and increased serum HDL-cholesterol.

Avocado may modify the structure of the HDL lipoprotein by increasing paraoxonase1 (PON1) enzyme activity. The cardioprotector effect of HDL-cholesterol is in part due of PON1 activity, which is responsible for the hydrolysis of lipid hydroperoxides (products of the lipid oxidation) [33] [34]. About LDL-cholesterol particles, a randomized controlled trial was conducted among 45 overweight/obese participants submitted to a 2-week run-in diet (Average American Diet, AAD) and later allocated to three distinct diets: 1) LF—low fat diet (24% of daily total calories from fats); 2) MF—moderate fat diet (34% of daily total calories from fats); or 3) AV—avocado diet (~140 g of Hass avocado a day 13 g of MUFA). The AV diet group showed higher reduction of LDL and non-HDL cholesterol when compared to the other diets, and had reduced values of total LDL-cholesterol particles (LDL-P -80.1 nmol/l,  $p = 0.0001$ ), subclasses of LDL-cholesterol (LDL<sub>3+4</sub>-4.1 mg/dl,  $p = 0.04$ ) and LDL/HDL ratio (-6.6%,  $p < 0.0001$ ). AV and MF diets reduced apolipoprotein B-100 (ApoB) levels and the relationship between ApoB/ApoA-I decreased in AV diet group, without weight changes [35].

### 1.3. Avocado and Nutritional Status

The pathophysiological processes linking obesity to atherosclerosis and cardiovascular disease clearly involve a chronic inflammatory state [36], which interact with other factors such as ectopic fat, insulin resistance and HTN. For weight management and prevention of cardiovascular disease, international dietary recommendations regarding MUFA ingestion vary from 12% to 25% of total daily calories [37].

Popularly, avocado is known by its high caloric value being “a fruit whose consumption should be contraindicated in diets for weight loss.” In fact, few studies have evaluated the avocado in the weight loss setting.

Pieterse *et al.* conducted a clinical trial among 61 overweight/obese subjects, who were randomly assigned in two groups (an isocaloric diet with or without 200 g/day of avocado). After 6 weeks individuals in both groups reduced their weight, Body Mass Index (BMI) and body fat percentage, without significant differences according to group. In other study for weight maintenance, 131 subjects were allocated for three diets: 1) a moderate fat diet (34% to 45% of total daily calories from fats, including 100 g/day of avocado); 2) a low fat diet (20% to 30% of total daily calories from fats); or 3) a control diet (35% of total calories from fats). After 6 months all groups showed significant increase body weight, without differences between them [38]. In general, avocado consumers had a higher daily consumption of fruits and vegetables, essential foods included in a healthy diet for reduction or maintenance of the body weight [39] [40].

### 1.4. Avocado and Blood Pressure

Hypertension is the leading cause of worldwide mortality and is responsible for approximately 40% of deaths from cardiovascular disease, chronic kidney disease (CKD) and type 2 diabetes mellitus (T2DM) [41]. Beyond weight reduction, the adoption of a DASH diet is a part of the non-pharmacological treatment for higher levels of blood pressure (BP) (systolic and diastolic BP > 140/90 mmHg) [42].

The high content of potassium and lutein in the avocado may improve the BP values by controlling oxidative stress and inflammation<sup>5</sup>. In addition, diets rich in MUFA may improve systolic and diastolic BP levels when compared to diets with low content of MUFA [43].

Experimental studies have evaluated the potential hypotensive effect attributed to the aqueous extract of the avocado leaves [44]-[46]. Ojewola *et al.* [44] showed that the vascular dilation in consequence to the aqueous extract intake was the responsible for an anti-hypertensive effect of the avocado in rats. Authors suggest that different phytochemical components in the avocado's extract caused this effect. In other study, the aqueous extract from avocado seeds used as treatment for HTN in rats reduced BP levels and also improved the lipid profile [47]. In humans, subjects with overweight and moderate obesity who received a restricted caloric diet supplemented with 200 g of avocado a day do not had their BP levels reduced after the intervention [38].

## 2. Conclusions

The consumption of avocado seems to be related to cardiometabolic health by preventing traditional risk factors such as dyslipidemia, glycemic control and hypertension (Table 2). Despite all beneficial effects of avocado,

**Table 2.** Summary of clinical trials regarding avocado consumption and cardiovascular risk factors.

Author	Year	No. Participants	Trial Design	Consumption /follow-up	Intervention	Outcome (s)
Grant <i>et al.</i> [31]	1960	16	Crossover	Daily, 4 weeks/phase	Avocado addition of 0.5 - 1.5 units	↓ total cholesterol
Colquhoun <i>et al.</i> [21]	1992	15	Crossover	Daily, 1 week/phase	Diet rich in MUFAs supplemented with 300 g avocado vs. diet rich in complex carbohydrates	↓ total cholesterol, LDL and triglycerides, ↑ HDL concentration
Carranza <i>et al.</i> [32]	1995	16	Crossover	Daily, 4 week/phase	Diet rich in avocados (75% of total fat), vs. diet low in saturated fat	↓ total cholesterol and LDL, ↑ increased HDL concentration
Pieterse <i>et al.</i> [38]	2005	61	Parallel	Daily, 6 weeks	Hypocaloric diet with inclusion of 200 g of avocado or hypocaloric diet	No difference regarding weight reduction and blood pressure
Sloth <i>et al.</i> [39]	2009	131	Parallel	Daily, 6 months	Moderate fat diet with 100 g avocado or low-fat diet or control diet	No difference regarding weight reduction
Wien <i>et al.</i> [20]	2013	26	Crossover	1 meal, postmeal measurement	Adding avocado 68g at lunch	↑ satiety and ↓ of fasting insulin in postprandial period of 3 h
Wang <i>et al.</i> [35]	2015	45	Crossover	Daily, 5 weeks/phase	Low-fat diet, moderate fat diet or moderate fat diet with addition of 136 g of avocado per day	↓ LDL and weight maintenance

many studies were made in animal models and there results should be interpreted with caution. It is suggested that further studies must be designed among humans, in order to evaluate and to confirm the benefits of avocado.

## References

- [1] Montenegro, H.W.S. (1951) The Culture of Avocado. Melhoramentos, São Paulo, 11:102.
- [2] Maranca, G. (1980) Commercial Fruit Growing: Mango and Avocado. Nobel, São Paulo, 81-133.
- [3] Koller, O.C. (1992) Abacaticultura. UFRGS, Porto Alegre, 138.
- [4] Canto, W.L., Santos L.C. and Travaglini, M.M.E. (1978) Avocado: The Crop to Processing and Marketing. Série frutas tropicais-1. ITAL, Campinas, 212.
- [5] Dreher, M.L. and Davenport, A.J. (2013) Hass Avocado Composition and Potential Health Effects. *Critical Reviews in Food Science and Nutrition*, **53**, 738-750. <http://dx.doi.org/10.1080/10408398.2011.556759>
- [6] Naveh, E., Werman, M.J., Sabo, E. and Neeman, I. (2002) Defatted Avocado Pulp Reduces Body Weight and Total Hepatic Fat But Increases Plasma Cholesterol in Male Rats Fed Diets with Cholesterol. *The Journal of Nutrition*, **132**, 2015-2018.
- [7] USDA (U.S. Department of Agriculture) (2011) Avocado, Almond, Pistachio and Walnut Composition. Nutrient Data Laboratory. USDA National Nutrient Database for Standard Reference, Release 24. U.S. Department of Agriculture. Washington, DC.
- [8] Unlu, N., Bohn, T., Clinton, S.K. and Schwartz, S.J. (2005) Carotenoid Absorption from Salad and Salsa by Humans Is Enhanced by the Addition of Avocado or Avocado Oil. *The Journal of Nutrition*, **135**, 431-436.
- [9] Wang, L., Bordi, P.L., Fleming, J.A., Hill, A.M. and Kris-Etherton, P.M. (2015) Effect of a Moderate Fat Diet with and without Avocados on Lipoprotein Particle Number, Size and Subclasses in Overweight and Obese Adults: A Randomized, Controlled Trial. *Journal of the American Heart Association*, **4**, e001355. <http://dx.doi.org/10.1161/jaha.114.001355>
- [10] Fulgoni 3rd, V.L., Dreher, M. and Davenport, A.J. (2013) Avocado Consumption Is Associated with Better Diet Quality and Nutrient Intake, and Lower Metabolic Syndrome Risk in US Adults: Results from the National Health and Nutrition Examination Survey (NHANES) 2001-2008. *The Journal of Nutrition*, **12**, 1. <http://dx.doi.org/10.1186/1475-2891-12-1>
- [11] Nicholson, G. and Hall, G.M. (2011) Diabetes Mellitus: New Drugs for a New Epidemic. *British Journal of Anaesthesia*, **107**, 65-73. <http://dx.doi.org/10.1093/bja/aer120>
- [12] Tangvarasittichai, S. (2015) Oxidative Stress, Insulin Resistance, Dyslipidemia and Type 2 Diabetes Mellitus. *World Journal of Diabetes*, **6**, 456-480. <http://dx.doi.org/10.4239/wjd.v6.i3.456>



- [13] American Diabetes Association (2014) Standards of Medical Care in Diabetes. *Diabetes Care*, **37**, 14-80.
- [14] Alhassan, A.J., Sule, M.S., Atiku, M.K., Wudil, A.M., Abubakar, H. and Mohammed, S.A. (2012) Effects of Aqueous Avocado Pear (*Persea americana*) Seed Extract on Alloxan Induced Diabetes Rats. *Greener Journal of Medical Sciences*, **2**, 5-11.
- [15] N'guessan, K., Amoikon, K.E. and Soro, D. (2009) Effect of Aqueous Extract of *Persea americana* Seeds on the Glycemia of Diabetic Rabbits. *European Journal of Scientific Research*, **26**, 376-385.
- [16] Ezejiolor, A.N., Okorie, A. and Orisakwe, O.E. (2013) Hypoglycaemic and Tissue-Protective Effects of the Aqueous Extract of *Persea americana* Seeds on Alloxan-Induced Albino Rats. *The Malaysian Journal of Medical Sciences*, **20**, 31-39.
- [17] Oboh, G., Isaac, A.T., Akinyemi, A.J. and Ajani, R.A. (2014) Inhibition of Key Enzymes Linked to Type 2 Diabetes and Sodium Nitroprusside Induced Lipid Peroxidation in Rats' Pancreas by Phenolic Extracts of Avocado Pear Leaves and Fruit. *International Journal of Biomedical Sciences*, **10**, 208-216.
- [18] Rao, U.S. and Adinew, B. (2011) Remnant B-Cell-Stimulative and Anti-Oxidative Effects of *Persea americana* Fruit Extract Studied in Rats Introduced into Streptozotocin-Induced Hyperglycaemic State. *African Journal of Traditional, Complementary and Alternative Medicines*, **8**, 210-217. <http://dx.doi.org/10.4314/ajtcam.v8i3.65277>
- [19] Ros, E. (2003) Dietary Cis-Monounsaturated Fatty Acids and Metabolic Control in Type 2 Diabetes. *The American Journal of Clinical Nutrition*, **78**, 617-625.
- [20] Wien, M., Haddad, E., Oda, K. and Sabaté, J. (2013) A Randomized 3 × 3 Crossover Study to Evaluate the Effect of Hass Avocado Intake on Post-Ingestive Satiety, Glucose and Insulin Levels, and Subsequent Energy Intake in Overweight Adults. *Nutrition Journal*, **12**, 155. <http://dx.doi.org/10.1186/1475-2891-12-155>
- [21] Colquhoun, D.M., Moores, D., Somerset, S.M. and Humphries, J.A. (1992) Comparison of the Effects on Lipoproteins and Apolipoproteins of a Diet High in Monounsaturated Fatty Acids, Enriched with Avocado, and a High-Carbohydrate Diet. *The American Journal of Clinical Nutrition*, **56**, 671-677.
- [22] Pahua-Ramos, M.E., Garduño-Siciliano, L., Dorantes-Alvarez, L., Chamorro-Cevallos, G., Herrera-Martínez, J., Osorio-Esquivel, O., *et al.* (2014) Reduced-Calorie Avocado Paste Attenuates Metabolic Factors associated with a Hypercholesterolemic-High Fructose Diet in Rats. *Plant Foods for Human Nutrition*, **69**, 18-24. <http://dx.doi.org/10.1007/s11130-013-0395-4>
- [23] Mendis, S., Puska, P. and Norrving, B. (2011) Global Atlas on Cardiovascular Disease Prevention and Control. World Health Organization.
- [24] World Health Organization (2009) Global Health Risks: Mortality and Burden of Disease Attributable to Selected Major Risks. World Health Organization.
- [25] Eckel, R.H., Jakicic, J.M., Ard, J.D., de Jesus, J.M., Houston, M.N., Hubbard, V.S., *et al.* (2013) AHA/ACC Guideline on Life Style Management to Reduce Cardiovascular Risk: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*, **129**, S76-S99. <http://dx.doi.org/10.1161/01.cir.0000437740.48606.d1>
- [26] Xavier, H.T., Izar, M.C., Faria Neto, J.R., Assad, M.H., Rocha, V.Z., Sposito, A.C., *et al.* (2013) V Brazilian Guideline of Dyslipidemia and Prevention of Atherosclerosis. *Arquivos Brasileiros de Cardiologia*, **101**, 1-20. <http://dx.doi.org/10.5935/abc.2013S010>
- [27] Santos, R.D., Gagliardi, A.C.M., Xavier, H.T., Magnoni, C.D., Cassani, R. and Lottenberg, A.M. (2013) Guidelines on the Consumption of Fats and Cardiovascular Health. *Arquivos Brasileiros de Cardiologia*, **100**, 1-40.
- [28] Sposito, A.C., Caramelli, B., Fonseca, F.A.H., Bertolami, M.C., Afiune Neto, A., Souza, A.D., *et al.* (2007) IV Brazilian Guideline of Dyslipidemia and Prevention of Atherosclerosis: Department of Atherosclerosis of the Brazilian Society of Cardiology. *Arquivos Brasileiros de Cardiologia*, **88**, 2-19. <http://dx.doi.org/10.1590/S0066-782X2007000700002>
- [29] Estruch, R., Ros, E., Salas-Salvado, J., Covas, M.I., Corella, D., Arós, F., *et al.* (2013) Primary Prevention of Cardiovascular Disease with Mediterranean Diet. *The New England Journal of Medicine*, **368**, 1279-1290. <http://dx.doi.org/10.1056/NEJMoa1200303>
- [30] Salgado, J.M., Bin, C., Mansi, D.N. and Souza, A. (2008) Effects of Avocado (*Persea americana* Mill) Variety Hass in Blood Lipids of Hypercholesterolemic Rats. *Ciência e Tecnologia dos Alimentos*, **28**, 922-928. <http://dx.doi.org/10.1590/S0101-20612008000400025>
- [31] Grant, W.C. (1960) Influence of Avocados on Serum Cholesterol. *Proceedings of the Society for Experimental Biology and Medicine*, **104**, 45-47. <http://dx.doi.org/10.3181/00379727-104-25722>
- [32] Carranza, J., Alvizouri, M., Alvarado, M.R., Chávez, F., Gómez, M. and Herrera, J.E. (1995) Effects of Avocado on the Level of Blood Lipids in Patients with Phenotype II and IV Dyslipidemias. *Archivos del Instituto de Cardiología de México*, **65**, 342-348.

- [33] Pérez Méndez, O. and García Hernández, L. (2007) High-Density Lipoproteins (HDL) Size and Composition Are Modified in the Rat by a Diet Supplemented with “Hass” Avocado (*Persea americana* Miller). *Archivos del Instituto de Cardiología de México*, **77**, 17-24.
- [34] Boshtam, M., Razavi, A.E., Pourfarzam, M., Ani, M., Naderi, G.A., Basati, G., *et al.* (2013) Serum Paraoxonase Activity Is Associated with Fatty Acid Composition of High Density Lipoprotein. *Diseases Markers*, **35**, 273-280. <http://dx.doi.org/10.1155/2013/612035>
- [35] Wang, L., Bordi, P.L., Fleming, J.A., Hill, A.M. and Kris-Etherton, P.M. (2015) Effect of a Moderate Fat Diet with and without Avocados on Lipoprotein Particle Number, Size and Subclasses in Overweight and Obese Adults: Randomized, Controlled Trial. *Journal of the American Heart Association*, **4**, e001355. <http://dx.doi.org/10.1161/jaha.114.001355>
- [36] Bastien, M., Poirier, P., Leimeux, I. and Després, J.P. (2014) Overview of Epidemiology and Contribution of Obesity to Cardiovascular Disease. *Progress in Cardiovascular Disease*, **56**, 369-381. <http://dx.doi.org/10.1016/j.pcad.2013.10.016>
- [37] Schwingshackl, L. and Hoffmann, G. (2012) Monounsaturated Fatty Acids and Risk of Cardiovascular Disease: Synopsis of the Evidence Available from Systematic Reviews and Meta-Analyses. *Nutrients*, **4**, 1989-2007. <http://dx.doi.org/10.3390/nu4121989>
- [38] Pieterse, E., Jerling, J.C., Oosthuizen, W., Kruger, H.S., Hanekom, S.M., Smuts, C.M., *et al.* (2005) Substitution of High Monounsaturated Fatty Acid Avocado for Mixed Dietary Fats during an Energy-Restricted Diet: Effects on Weight Loss, Serum Lipids, Fibrinogen, and Vascular Function. *Nutrition*, **21**, 67-75. <http://dx.doi.org/10.1016/j.nut.2004.09.010>
- [39] Sloth, B., Due, A., Larsen, T.M., Holst, J.J., Heding, A. and Astrup, A. (2009) The Effect of a High-MUFA, Low-Glycaemic Index Diet and a Low-Fat Diet on Appetite and Glucose Metabolism during a 6-Month Weight Maintenance Period. *British Journal of Nutrition*, **101**, 1846-1858. <http://dx.doi.org/10.1017/S0007114508137710>
- [40] Champagne, C.M., Broyles, S.T., Moran, L.D., Cash, K.C., Levy, E.J., Lin, P.-H., *et al.* (2011) Dietary Intakes Associated with Successful Weight Loss and Maintenance during the Weight Loss Maintenance Trial. *Journal of the American Dietetic Association*, **111**, 1826-1835. <http://dx.doi.org/10.1016/j.jada.2011.09.014>
- [41] Schmidt, M.S., Duncan, B.B., Azevedo e Silva, G., Menezes, A.M., Monteiro, C.A., Barreto, S.M., *et al.* (2011) Chronic Non-Communicable Diseases in Brazil: Burden and Current Challenges. *Lancet*, **377**, 1949-1961. [http://dx.doi.org/10.1016/S0140-6736\(11\)60135-9](http://dx.doi.org/10.1016/S0140-6736(11)60135-9)
- [42] Mancia, G., Fagard, R., Narkiewicz, K., Redon, J., Zanchetti, A., Bohm, M., *et al.* (2013) ESH/ESC Guidelines for the Management of Arterial Hypertension: The Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *European Heart Journal*, **34**, 2159-2219. <http://dx.doi.org/10.1093/eurheartj/eh1151>
- [43] Schwingshackl, L., Strasser, B. and Hoffmann, G. (2011) Effects of Monounsaturated Fatty Acids on Cardiovascular Risk Factors: A Systematic Review and Meta-Analysis. *Annals of Nutrition and Metabolism*, **59**, 176-186. <http://dx.doi.org/10.1159/000334071>
- [44] Ojewole, J.A., Kamadyaapa, D.R., Gondwe, M.M., Moodley, K. and Musabayane, C.T. (2007) Cardiovascular Effects of *Persea americana* Mill (Lauraceae) (Avocado) Aqueous Leaf Extract in Experimental Animals. *Cardiovascular Journal of Africa*, **18**, 69-76.
- [45] Adeboye, J.O., Fajonyomi, M.O., Makinde, J.M. and Taiwo, O.B. (1999) A Preliminary Study on the Hypotensive Activity of *Persea americana* Leaf Extracts in Anaesthetized, Normotensive Rats. *Fitoterapia*, **70**, 15-20. [http://dx.doi.org/10.1016/S0367-326X\(98\)00015-X](http://dx.doi.org/10.1016/S0367-326X(98)00015-X)
- [46] Owolabi, M.A., Jaja, S.I. and Coker, H.A.B. (2005) Vasorelaxant Action of Aqueous Extract of the Leaves of *Persea americana* on Isolated Thoracic Rat Aorta. *Fitoterapia*, **76**, 567-573. <http://dx.doi.org/10.1016/j.fitote.2005.04.020>
- [47] Imafidon, K.E. and Amaechina, F.C. (2010) Effects of Aqueous Seed Extract of *Persea americana* Mill. (Avocado) on Blood Pressure and Lipid Profile in Hypertensive Rats. *Advances in Biological Research*, **4**, 116-121.